



SCHOOL OF MATHEMATICS AND STATISTICS

Spring Semester
2013–2014

MAS6450 Waves and Magnetohydrodynamics

2 hours

Answer all four questions.

- 1 (i) Consider a magnetic field with components

$$B_x(x, z) = \frac{\partial\psi}{\partial z}, \quad B_y = B_y(x, z), \quad B_z(x, z) = -\frac{\partial\psi}{\partial x},$$

where $\psi = \psi(x, z)$.

- (a) Show that $\nabla \cdot \mathbf{B} = 0$. (2 marks)
- (b) Show that $\mathbf{B} \cdot \nabla\psi = 0$ and that projections of field lines in the xz -plane are given by $\psi = \text{constant}$. (6 marks)
- (c) Show that if

$$\mathbf{J} \times \mathbf{B} = \mathbf{0},$$

where \mathbf{J} is the current density, then

$$B_y = B_y(\psi)$$

and ψ satisfies

$$\frac{\partial^2\psi}{\partial x^2} + \frac{\partial^2\psi}{\partial z^2} + B_y \frac{dB_y}{d\psi} = 0.$$

(12 marks)

- (ii) Calculate the approximate timescale (in years) for the decay of the interstellar magnetic field given the parameters $L = 3 \times 10^{16}$ m and $\eta = 3.6 \times 10^6$ cm² s⁻¹. (5 marks)

- 2 (i) State the relationship between Lagrangian and Eulerian perturbations. Write the Lagrangian density perturbation in terms of the Eulerian perturbation. (4 marks)

2 (continued)

(ii) Consider

$$\mathbf{B} = B_0(y/a, x/a, 0).$$

Sketch the field lines showing the direction. *(4 marks)*

Calculate the magnetic tension and magnetic pressure forces. Comment on the direction of each. Give comments about the Lorentz force. *(7 marks)*

(iii) Prove that

$$\frac{\partial \hat{\boldsymbol{\theta}}}{\partial \theta} = -\hat{\mathbf{r}},$$

where r and θ are two-dimensional polar coordinates with $\hat{\mathbf{r}}$ and $\hat{\boldsymbol{\theta}}$ being the corresponding unit vectors. *(3 marks)*

(iv) Prove, using (iii), that for magnetic fields of the form $\mathbf{B} = B_\theta(r)\hat{\boldsymbol{\theta}}$,

$$(\mathbf{B} \cdot \nabla)\mathbf{B} = -\frac{B_\theta^2}{r}\hat{\mathbf{r}}.$$

(3 marks)

3 (i) (a) Ignoring viscosity, gravity and diffusivity, derive the linearised induction equation and equation of motion for adiabatic, ideal perturbation of the form $\sim e^{i\mathbf{k}\cdot\mathbf{r}-\omega t}$, about a static and uniform equilibrium state with magnetic force alone. *(6 marks)*

[You may use $\nabla \times (\mathbf{A} \times \mathbf{B}) = (\mathbf{B} \cdot \nabla)\mathbf{A} - (\mathbf{A} \cdot \nabla)\mathbf{B} + (\nabla \cdot \mathbf{B})\mathbf{A} - (\nabla \cdot \mathbf{A})\mathbf{B}$ and $\mathbf{A} \times (\mathbf{B} \times \mathbf{C}) = (\mathbf{A} \cdot \mathbf{C})\mathbf{B} - (\mathbf{A} \cdot \mathbf{B})\mathbf{C}$].

(b) Derive the dispersion relations for Alfvén and compressional Alfvén waves using linearised MHD equations obtained in (i). *(8 marks)*

(c) State one property of each:

(a) Alfvén waves

(b) compressional Alfvén waves. *(2 marks)*

(ii) (a) Write the induction equation for a perfectly conducting medium. Given a velocity field $\mathbf{v} = (yz, -xz, 0)$ and the initial condition on the magnetic field

$$\mathbf{B}(\mathbf{x}, 0) = (x, -y, 0),$$

find $\mathbf{B}(\mathbf{x}, t)$ by obtaining the Lagrangian coordinates corresponding to \mathbf{v} and applying the Cauchy solution. *(19 marks)*

3 (continued)

Hint: Use an initial condition $\mathbf{x}(0) = (a_1, a_2, a_3)$ and also make use of $\mathbf{x}(0)$ for the initial magnetic field $\mathbf{B}(\mathbf{x}, 0) = \mathbf{B}_0$.

(b) Having obtained $\mathbf{B}(\mathbf{x}, t)$ in (iv), verify by direct substitution that it is indeed the solution of the induction equation. **(4 marks)**

4 (i) Consider a cartesian coordinate system where the magnetic and velocity fields are independent of y , i.e.

$$\mathbf{B} = \left(-\frac{\partial A}{\partial z}, B, \frac{\partial A}{\partial x} \right),$$

$$\mathbf{v} = \left(-\frac{\partial \psi}{\partial z}, v_y, \frac{\partial \psi}{\partial x} \right),$$

where A and ψ are scalar potential functions for magnetic and flow fields, respectively. Using the magnetic induction equation, the equations for the evolution of B and A can be given as

$$\frac{\partial B}{\partial t} + \left(\frac{\partial \psi}{\partial x} \frac{\partial B}{\partial z} - \frac{\partial \psi}{\partial z} \frac{\partial B}{\partial x} \right) = \left(\frac{\partial A}{\partial x} \frac{\partial v_y}{\partial z} - \frac{\partial A}{\partial z} \frac{\partial v_y}{\partial x} \right) - \left(\nabla \cdot (\alpha \nabla A) - \eta \nabla^2 B \right), \quad (\text{I})$$

$$\frac{\partial A}{\partial t} + \left(\frac{\partial \psi}{\partial x} \frac{\partial A}{\partial z} - \frac{\partial \psi}{\partial z} \frac{\partial A}{\partial x} \right) = \alpha B + \eta \left[\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right] A. \quad (\text{II})$$

Simplify the above equations (I and II) by setting $\psi = 0$, $v_y = Qz$ and ignoring the term containing α in the equation for the evolution of B .

(2 marks)

(ii) Assuming solutions of the form $(A, B) = (\tilde{A}, \tilde{B})e^{\sigma t - i(k_x x + k_z z)}$ in the simplified equations obtained in (i), show that the following condition is satisfied for the solutions to exist

$$(\sigma + \eta \kappa^2)^2 = -i k_x \alpha Q,$$

where $\kappa^2 = k_x^2 + k_z^2$.

(6 marks)

(iii) In the equations for the evolution of B and A in (i), assume $\psi = v_y = 0$ but retain the term containing α to be a constant. Substitute the same solutions of the form $(A, B) = (\tilde{A}, \tilde{B})e^{\sigma t - i(k_x x + k_z z)}$ and show that now the solution satisfies the relation

$$\sigma = \pm \alpha \kappa - \eta \kappa^2.$$

(7 marks)

End of Question Paper