



The  
University  
Of  
Sheffield.

**MAS6352**

**SCHOOL OF MATHEMATICS AND STATISTICS**

**Spring Semester  
2014–2015**

**MAS6352 Analysis II**

**2 hours**

*Full marks may be obtained by complete answers to three questions. All answers will be marked, but credit will be given only for the best three answers. Total marks 99.*

1 (i) (a) Let  $S$  be a set. List the properties that  $\Sigma$  must have for  $(S, \Sigma)$  to be a measurable space. (3 marks)

(b) Explain what is meant by a measure on  $(S, \Sigma)$ . (2 marks)

(ii) Let  $\alpha$  and  $\beta$  be positive real numbers, and  $m_1$  and  $m_2$  be measures on  $(S, \Sigma)$ .

(a) Show that  $\alpha m_1 + \beta m_2$  is a measure on  $(S, \Sigma)$ , where for each  $A \in \Sigma$ ,

$$(\alpha m_1 + \beta m_2)(A) = \alpha m_1(A) + \beta m_2(A).$$

(4 marks)

(b) Suppose that  $m_1$  and  $m_2$  are both probability measures. Give a condition on  $\alpha$  and  $\beta$  so that  $\alpha m_1 + \beta m_2$  is also a probability measure.

(3 marks)

(iii) (a) If  $A, B, C \in \Sigma$  with  $C \subset B \subset A$ , deduce that

$$m(B \cap (A - C)) = m(B) - m(C).$$

(3 marks)

(b) Calculate the Lebesgue measure of the Borel subset  $E$  of  $\mathbb{R}$  defined by

$$E = \left[0, \frac{1}{2}\right] \cap \left\{ [0, 1] - \bigcup_{n=1}^{\infty} \left(\frac{1}{7^{n+1}}, \frac{1}{7^n}\right) \right\}.$$

(5 marks)

(iv) Let  $(S, \Sigma)$  be a measurable space and  $(f_n)$  be a sequence of bounded measurable functions from  $S$  to  $\mathbb{R}$ .

(a) Define  $\liminf_{n \rightarrow \infty} f_n$  and  $\limsup_{n \rightarrow \infty} f_n$  and explain why these functions are measurable. (5 marks)

(b) Suppose that  $\liminf_{n \rightarrow \infty} f_n = \limsup_{n \rightarrow \infty} f_n$  almost everywhere. What can you say about the existence and measurability of  $\lim_{n \rightarrow \infty} f_n$ ? (4 marks)

(c) Define a sequence of functions  $(f_n)$  from  $\mathbb{R}$  to  $\mathbb{R}$  by

$$f_n(x) = \begin{cases} \left(1 + \frac{x}{n}\right) & \text{if } x \in (0, 1) \\ 0 & \text{if } x \notin (0, 1). \end{cases}$$

Is each  $f_n$  bounded and measurable? Does  $\lim_{n \rightarrow \infty} f_n(x)$  exist for all  $x \in \mathbb{R}$ , and if so, does the limit give rise to a measurable function? (4 marks)

- 2 (i) Let  $(S, \Sigma)$  be a measurable space. Explain what is meant by both a  $\pi$ -system, and a  $\lambda$ -system. State Dynkin's  $\pi - \lambda$  lemma. **(6 marks)**

From now on let  $(S_1, \Sigma_1)$  and  $(S_2, \Sigma_2)$  be measure spaces. If  $E \in \Sigma_1 \otimes \Sigma_2$  and  $y \in S_2$ , let  $E_y$  be the  $y$ -slice of  $E$  defined by

$$E_y = \{x \in S_1, (x, y) \in E\}.$$

You may freely use the facts that

$$(E^c)_y = (E_y)^c, \quad (E - F)_y = E_y - F_y, \quad \left( \bigcup_{n=1}^{\infty} E_n \right)_y = \bigcup_{n=1}^{\infty} (E_n)_y,$$

and that  $E_y \in \Sigma_1$ .

- (ii) If  $f : S_1 \times S_2 \rightarrow \mathbb{R}$  is measurable, show that  $f_y : S_1 \rightarrow \mathbb{R}$  is also measurable, where  $f_y(x) = f(x, y)$  for all  $x \in S_1$ . **(4 marks)**
- (iii) Let  $m_1$  and  $m_2$  be finite measures on  $(S_1, \Sigma_1)$  and  $(S_2, \Sigma_2)$ , respectively, and for each  $E \in \Sigma_1 \otimes \Sigma_2$ , define  $\psi_E : S_2 \rightarrow \mathbb{R}$  by

$$\psi_E(y) = m_1(E_y).$$

In order to define the product measure  $(m_1 \times m_2)(E) = \int_{S_2} m_1(E_y) dm_2(y)$ , we must first show that  $\psi_E$  is measurable. Define

$$\mathcal{S} = \{E \in \Sigma_1 \otimes \Sigma_2; \psi_E \text{ is measurable}\}.$$

- (a) Show that  $\mathcal{S}$  is a  $\lambda$ -system. **(10 marks)**
- (b) Show that for all  $A \in \Sigma_1, B \in \Sigma_2$ , the product set  $A \times B \in \mathcal{S}$ . **(3 marks)**
- (c) Show that  $\{A \times B; A \in \Sigma_1, B \in \Sigma_2\}$  forms a  $\pi$ -system. **(2 marks)**
- (d) Deduce that  $\psi_E$  is measurable. **(2 marks)**
- (iv) (a) If  $g : S_2 \rightarrow \mathbb{R}$  is measurable and  $A \in \Sigma_1$ , deduce that  $\tilde{g}_A : S_1 \times S_2 \rightarrow \mathbb{R}$  is measurable, where  $\tilde{g}_A(x, y) = \mathbf{1}_A(x)g(y)$ , for all  $x \in S_1, y \in S_2$ . **(2 marks)**
- (b) Suppose that  $f : S_1 \rightarrow \mathbb{R}$  is measurable. Use the result of (a) to show that  $h : S_1 \times S_2 \rightarrow \mathbb{R}$  is measurable, where  $h(x, y) = f(x)g(y)$ , for all  $x \in S_1, y \in S_2$ . **(4 marks)**

**3** Throughout this question  $(S, \Sigma, m)$  is a measure space.

- (i) State the *monotone convergence theorem*, and use it to prove *Fatou's lemma*: if  $(f_n)$  is a sequence of non-negative measurable functions then

$$\liminf_{n \rightarrow \infty} \int_S f_n dm \geq \int_S \liminf_{n \rightarrow \infty} f_n dm.$$

*(8 marks)*

- (ii) Suppose that the sequence  $(f_n)$  of (i) converges pointwise to  $f$ . Deduce that

$$\limsup_{n \rightarrow \infty} \int_S f_n dm \geq \int_S f dm.$$

*(3 marks)*

- (iii) Let  $f : S \rightarrow \mathbb{R}$  be a non-negative measurable function and define a sequence  $(f_n)$  of functions by  $f_n(x) = \min\{f(x), n\}$  for all  $x \in S, n \in \mathbb{N}$ . Deduce that

$$\lim_{n \rightarrow \infty} \int_S f_n dm = \int_S f dm.$$

*(5 marks)*

- (iv) State the *dominated convergence theorem*, and use it to find

(a)

$$\lim_{n \rightarrow \infty} \int_0^\infty \frac{1 - e^{-nx}}{(1 + x^2)(1 - e^{-2nx})} dx.$$

*(9 marks)*

(b)

$$\lim_{n \rightarrow \infty} \int_0^1 \frac{1 + nx}{(1 + x)^n} dx.$$

*(8 marks)*

4 Throughout this question,  $(\Omega, \mathcal{F}, P)$  is a probability space.

- (i) Let  $(A_n)$  be a sequence of events in  $\mathcal{F}$ . Define the events  $\liminf_{n \rightarrow \infty} A_n$  and  $\limsup_{n \rightarrow \infty} A_n$ , and deduce that

$$P\left(\liminf_{n \rightarrow \infty} A_n^c\right) = 1 - P\left(\limsup_{n \rightarrow \infty} A_n\right)$$

**(6 marks)**

- (ii) State both parts of the *Borel–Cantelli lemma*, and prove the part that does not require an independence assumption. **(6 marks)**
- (iii) Consider a sequence of (independent) rolls of a fair die. Deduce that the run 1412 appears infinitely often. **(7 marks)**
- (iv) Let  $(X_n)$  be a sequence of random variables which is such that for all  $\epsilon > 0$ ,  $P\left(\limsup_{n \rightarrow \infty} (|X_n| \geq \epsilon)\right) = 0$ . What can you say about the asymptotic behaviour of  $X_n$  as  $n \rightarrow \infty$ ? **(6 marks)**
- (v) The characteristic function of a random variable  $X$  is the mapping  $\Phi_X : \mathbb{R} \rightarrow \mathbb{C}$  defined by  $\Phi_X(u) = \mathbb{E}(e^{iuX})$ , for all  $u \in \mathbb{R}$ .
- (a) Show that the mapping  $u \rightarrow \Phi_X(u)$  is continuous from  $\mathbb{R}$  to  $\mathbb{C}$ . **(3 marks)**
- (b) Write down the characteristic functions for the following random variables, (I)  $X \sim N(a, \sigma^2)$ , (II)  $X = a$  (a.s.), where  $a \in \mathbb{R}$ ,  $\sigma > 0$ . **(2 marks)**
- (c) What can you say about the asymptotic behaviour of the sequence  $(X_n)$  of random variables, where  $X_n \sim N(a, 1/n)$ ? **(3 marks)**

**End of Question Paper**