



The
University
Of
Sheffield.

MAS153/MAS159

SCHOOL OF MATHEMATICS AND STATISTICS

**Spring Semester
2018–2019**

**Mathematics (Materials)
Mathematics For Chemists**

3 hours

All questions are compulsory. The marks awarded to each question or section of question are shown in italics. Total Marks 100

- 1 Find all solutions of the equation

$$2 \cos^2 x - 3 \cos x + \sin^2 x + 1 = 0. \quad (5 \text{ marks})$$

- 2 Solve the equation

$$\ln[(x-2)^4] = 2 \ln(x-1) + \ln[(x+1)^2]. \quad (6 \text{ marks})$$

- 3 Identify the number d such that

$$5^d = \frac{5 \times (21)^{\frac{2}{3}} \times 10}{(27)^{\frac{2}{9}} \times 2 \times 7^{\frac{2}{3}}}. \quad (4 \text{ marks})$$

- 4 Simplify

$$\frac{(8x^3y^{\frac{3}{2}}z^6)^{\frac{2}{3}}}{2x^2(y^3\sqrt{z})^2}. \quad (4 \text{ marks})$$

- 5 Factorise $3x^2 + 9x - 12$.

(3 marks)

6 Verify the following identity

$$(\tan^2 x + 1)^2(1 - \sin^2 x)^2 = 1. \quad (5 \text{ marks})$$

7 Solve the following equation for real x :

$$25^x - 6 \times 5^x + 5 = 0. \quad (5 \text{ marks})$$

8 Let $y = e^{\sin(x)+x^2}$. Find $\frac{dy}{dx}$ and $\frac{d^2y}{dx^2}$. (5 marks)

9 (i) Showing your working clearly, find the coefficient of x^3 in the expansion of $(1 + 2x)^{10}$. (2 marks)

(ii) Use the binomial theorem to evaluate

$$\lim_{x \rightarrow \infty} (\sqrt{x^2 + 4x + 5} - x - 3). \quad (4 \text{ marks})$$

10 (i) Find the equation of the lines passing through the following pairs of points:

(a) $(0, 0, -1)$ and $(1, 5, 1)$;

(b) $(1, 1, 1)$ and $(2, 1, 3)$;

(c) $(5, 3, 7)$ and $(9, 1, 17)$. (3 marks)

(ii) On which of the above lines (i.e. from part i), if any, do the following points lie:

(a) $\frac{1}{5}(1, 5, -3)$;

(b) $(1, 5, 1)$;

(c) $(3, 2, 3)$. (5 marks)

11 (i) Prove, using the exponential form,

$$\coth^2(x) - 1 = \operatorname{cosech}^2(x). \quad (4 \text{ marks})$$

(ii) Find the inverse of $\tanh(x)$. (4 marks)

12 Find

$$\int \frac{x^2 - x + 1}{x^3 - 2x^2 + x - 2} dx$$

(9 marks)

13 Find the Maclaurin Series for $\cos(2x^2)$, up to terms in x^6 .

(8 marks)

14 Using De Moivre's theorem find $\cos(4\theta)$ in terms of $\cos(\theta)$, and $\sin(4\theta)$ in terms of $\sin(\theta)$.

(12 marks)

15 A set of linear equations can be written as

$$A \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}.$$

where

$$A = \begin{bmatrix} 1 & 2 & 5 \\ 0 & \lambda & 3 \\ 1 & \lambda & 2 \end{bmatrix}.$$

Find λ such that there is a non-trivial solution. Using this value of λ find values of x, y and z which satisfy the equations.

(12 marks)

End of Question Paper

Formula Sheet for MAS153/MAS159 Examination

These results may be quoted without proof, unless proofs are asked for in the question.

Trigonometry

For any angles A and B

$$\sin^2 A + \cos^2 A = 1$$

$$\cos(A \pm B) = \cos A \cos B \mp \sin A \sin B$$

$$\sin(A \pm B) = \sin A \cos B \pm \cos A \sin B$$

$$\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$$

$$\sin 2A = 2 \sin A \cos A$$

$$\cos 2A = 2 \cos^2 A - 1 = 1 - 2 \sin^2 A$$

Coordinate Geometry

The acute angle α between lines with gradients m_1 and m_2 satisfies

$$\tan \alpha = \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right| \quad (m_1 m_2 \neq -1)$$

while the lines are perpendicular if $m_1 m_2 = -1$.

The equation of a circle centre (x_0, y_0) and radius a is $(x - x_0)^2 + (y - y_0)^2 = a^2$.

Hyperbolic Functions

$$\cosh^2 x - \sinh^2 x = 1$$

$$\operatorname{sech}^2 x + \tanh^2 x = 1$$

$$\cosh^2 x + \sinh^2 x = \cosh 2x$$

$$2 \sinh x \cosh x = \sinh 2x$$

$$\cosh^2 x = (1 + \cosh 2x)/2$$

$$\sinh^2 x = -(1 - \cosh 2x)/2$$

Differentiation

<u>Function</u> (y)	<u>Derivative</u> (dy/dx)
x^n	nx^{n-1}
$\sin ax$	$a \cos ax$
$\cos ax$	$-a \sin ax$
$\tan ax$	$a \sec^2 ax$
e^{ax}	ae^{ax}
$\ln(ax)$	$\frac{1}{x}$
$\ln f(x)$	$\frac{f'(x)}{f(x)}$
$\sinh x$	$\cosh x$
$\cosh x$	$\sinh x$
$\tanh x$	$\operatorname{sech}^2 x$
$\sin^{-1} x$	$\frac{1}{\sqrt{1-x^2}}$
$\cos^{-1} x$	$-\frac{1}{\sqrt{1-x^2}}$
$\tan^{-1} x$	$\frac{1}{1+x^2}$
$\sinh^{-1} x$	$\frac{1}{\sqrt{x^2+1}}$
$\cosh^{-1} x$	$\frac{1}{\sqrt{x^2-1}}$
$\tanh^{-1} x$	$\frac{1}{1-x^2}$

NB. It is assumed that x takes only those values for which the functions are defined.

For u and v functions of x , and with $u' = \frac{du}{dx}$, $v' = \frac{dv}{dx}$,

$$\frac{d}{dx}(uv) = uv' + vu',$$

while

$$\frac{d}{dx} \left(\frac{u}{v} \right) = \frac{vu' - uv'}{v^2}.$$

For $y = y(t)$, $t = t(x)$,

$$\frac{dy}{dx} = \frac{dy}{dt} \frac{dt}{dx}.$$

Integration

In the following table the constants of integration have been omitted.

<u>Function</u> $f(x)$	<u>Integral</u> $\int f(x) dx$
x^n	$\frac{x^{n+1}}{n+1} \quad n \neq -1$
ae^{ax}	e^{ax}
$\frac{1}{x}$	$\ln x $
$a \sin ax$	$-\cos ax$
$a \cos ax$	$\sin ax$
$a \tan ax$	$\ln \sec ax $
$\frac{1}{a^2 + x^2}$	$\frac{1}{a} \tan^{-1} \left(\frac{x}{a} \right)$
$\frac{1}{a^2 - x^2}$	$\frac{1}{a} \tanh^{-1} \left(\frac{x}{a} \right)$
$\frac{1}{\sqrt{a^2 - x^2}}$	$\sin^{-1} \left(\frac{x}{a} \right)$
$\frac{1}{\sqrt{x^2 + a^2}}$	$\sinh^{-1} \left(\frac{x}{a} \right)$
$\frac{1}{\sqrt{x^2 - a^2}}$	$\cosh^{-1} \left(\frac{x}{a} \right)$
$\frac{f'(x)}{f(x)}$	$\ln f(x) $

Integration by parts

$$\int uV dx = (\text{integral of } V) \times u - \int (\text{integral of } V) \times \frac{du}{dx} dx$$

or

$$\int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx.$$

Series

Binomial Theorem: $(1+x)^n = 1 + nx + \frac{n(n-1)}{2!}x^2 + \cdots + \binom{n}{r}x^r + \cdots$

$$\text{where } \binom{n}{r} = \frac{n(n-1)(n-2)\cdots(n-r+1)}{r!}$$

If n is a positive integer, the series terminates and is convergent for all x .

If n is not a positive integer, the series is infinite and converges for $|x| < 1$.

Taylor expansion of $f(x)$ about $x = a$ is

$$f(a) + (x-a)f^{(1)}(a) + \frac{(x-a)^2}{2!}f^{(2)}(a) + \cdots + \frac{(x-a)^n}{n!}f^{(n)}(a) + \cdots$$

Maclaurin expansion of $f(x)$ is

$$f(0) + xf^{(1)}(0) + \frac{x^2}{2!}f^{(2)}(0) + \cdots + \frac{x^n}{n!}f^{(n)}(0) + \cdots$$

Alternating Series Test

The series $a_1 - a_2 + a_3 - a_4 + \cdots$, where $a_1, a_2, a_3, a_4, \dots$ are all positive, converges if $a_1 > a_2 > a_3 > \cdots$ and $a_n \rightarrow 0$ as $n \rightarrow \infty$.

Ratio Test

If the series $a_1 + a_2 + a_3 + a_4 + \cdots$ satisfies

$$\lim_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| = \lambda,$$

then

1. if $\lambda > 1$, the series diverges,
2. if $\lambda < 1$, the series converges.

Vectors

If vectors \mathbf{a} and \mathbf{b} are given in Cartesian component form by $\mathbf{a} = (a_1, a_2, a_3)$ and $\mathbf{b} = (b_1, b_2, b_3)$, then

the scalar product $\mathbf{a} \cdot \mathbf{b}$ is given by

$$\mathbf{a} \cdot \mathbf{b} = a_1b_1 + a_2b_2 + a_3b_3$$

and the vector product $\mathbf{a} \times \mathbf{b}$ is given by

$$\mathbf{a} \times \mathbf{b} = \begin{vmatrix} \mathbf{i} & \mathbf{j} & \mathbf{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix} = (a_2b_3 - a_3b_2, a_3b_1 - a_1b_3, a_1b_2 - a_2b_1).$$

If a plane passes through a point with position vector \mathbf{a} , and is normal to the vector \mathbf{n} , then the equation of the plane is

$$\mathbf{r} \cdot \mathbf{n} = \mathbf{a} \cdot \mathbf{n},$$

where $\mathbf{r} = (x, y, z)$.